Review Article

Prolonged the Shelf Life of Different Foods using the Citrus By-Products as Antimicrobials: A Review Article

Al-Dalali S^{1,2*}, Zheng F¹ and Al-Farga A²

¹Beijing Advanced Innovation Center for Food Nutrition and Human Health, Beijing Technology and Business University, China

²Department of Food Science and Technology, Ibb University, Yemen

*Corresponding author: Sam Al-Dalali, Beijing Advanced Innovation Center for Food Nutrition and Human Health, Beijing Technology and Business University, China

Received: December 10, 2018; Accepted: January 22, 2019; Published: January 29, 2019

Abstract

A novel type of packaging that combines food packaging materials with antimicrobial substances to control microbial surface contamination of foods to enhance product microbial safety and to extend shelf-life is attracting interest in the packaging industry. In agreement with the current trend of giving value to natural and renewable resources, the use of natural antimicrobial compounds, particularly in food and biomedical applications, becomes very frequent. The direct addition of natural compounds to food is the most common method of application, even if numerous efforts have been made to find alternative solutions to the aim of avoiding undesirable inactivation. Dipping, spraying, and coating treatment of food with active solutions are currently applied to product prior to packaging as valid options.

Keywords: Antimicrobial compounds; Food preservation; Natural compounds; Food packaging; Natural antimicrobials

Prolonged Storage of Different Food by Using Citrus as Antimicrobial

At its known, the main factors that cause the deterioration in food quality and reduced their shelf life are bacterial growth and lipid oxidation [1,2]. So, there are demand for use chemical additives as a preservative to inhibit the spoilage of food by microorganism and lipid oxidation process. Nevertheless, nowadays the consumers become more aware about the defect of synthetic antimicrobial and antioxidants substances and their relationship about human health risk, which used for extend food shelf life [3]. So, the current tend to use the natural antimicrobial compounds for extend the shelf life of food [4]. Among of them, the antimicrobial and antioxidant that have been extracted from citrus, whether from peel, seed, pulp and essential oils and applied them in food products. Since its will known that the extracts consider as a high source of bio phytochemicals substances against microorganism, also they could be added in the food system to improve the quality and extend the shelf life [3].

A lot of phenolic compounds that have been reported in the citrus peels include flavonoids (flavonols, flavones, flavones and coumarins), phenolic acids and monoterpene hydrocarbons [5]. Many studies have been conducted by using essential oils extracts from citrus peels, the results exhibited that the antimicrobial activity against *Salmonella typhi, Bacillus cereus* and *Staphylococcus aureus* [6], also in fruit products [7], chicken products [8]. Extracted bio phytochemical from citrus can be added to food either directly or incorporated in packaging system [9]. A new preservation method that used to improve the quality of food products, it's called Edible Films and Coating (EFC) by added with natural antimicrobial to inhibited food spoilage and growth of pathogenic microorganism [10].

Meat and meat products

Meat and meat products are considered a good source of protein and other nutrients like vitamin and minerals. It's defined by the codex alimentarius [11] as a safe source and suitable for human consumption. Many of the factors that effect on the quality of meat such as pH, temperature, microbial growth and lipid oxidation ... etc. [12]. The deterioration that occur during the processing and storage of meat ref come back to three mechanisms: lipid oxidation, microbial spoilage and enzymatic autolysis [10].

Ethanol and ethyl acetate extraction of three type of citrus peel (kaffir lime, lime and pomelo) were studied in stored raw chicken drumettes at 4°c as antimicrobial and antioxidants for extension their shelf life. They were found to be more effective against pathogenic bacteria at level ranged from 0.4-50 mg/ml. also, they found that the treated chicken wing samples with ethyl acetate extracted from kaffir lime peel were lower than those of control sample in their values of total viable counts, 2-Thiobarbituric Acid Reactive Substances (TBARS) and increased the sensory properties during the period of storage (14 days) [13] (Table 1). In another study conducted by [14], investigate the effect of Citrus limon Essential Oil (CLEO) in situ against L. monocytogenes, as well as evaluated the physicochemical parameters (pH and TBARS) in minced beef meat model. They found that the CLEO was more effective against gram positive and negative bacteria at concentration varied from 0.039 to 1.25 mg/ml and 0.25 to 2.5 mg/ml respectively. Also, they assessed the CLEO as preservative substances in minced beef meet during storage at 4°c. The results showed that added 0.06 and 0.312 mg/g from CLEO were prohibit the contamination in pathogenic bacteria, particularly L. monocytogenes. Additionally, the values pf pH and TBARS were lower than those, which found in control meat. Ibrahim and Salem studied adding lemon grass and lime peel extracts (fresh and oil) to the prepared chicken patties during the storage for 9 days at 4°c, 0.25% of tested oil extracts ere an optimum concentration that have been effective against TBARS and lipid oxidation [15]. As well as, reduce the pH, aerobic plate count, mould and yeast in the treated chicken patties comparing the untreated (control). Additionally, the treated chicken patties were lower than in their content of Volatile Basic Nitrogen



(VBN) of control. On the other hand, they reported that addition the same concentration of lemo grass and lime peel extracts reduced the histamine, tyramine and putrescine formation in refrigerated chicken patties.

The extract of kinnow have been added as a source of antioxidant in salted chicken patties during storage. The results showed that the presence of salt alone improved the TBARS formation (114%), whereas adding both of the kinnow rind powder extract and salt at the same time led to reduced it to 90%. Also, the lipid oxidation was significantly reduced during the storage to 39% [16]. The effect of kinnow rind powder extract as a source of antioxidants in cooked goat's meat patties were investigated by Devatkal et al. [17]. The peel of citrus sinensis was dried and studied their effect on gastrointestinal microbiota and immune system of broiler chickens was conducted by Ebrahimi et al [18]. The results showed that the different amount of dried citrus sinensis peel was significantly affected on Escherichia coli in the caecum, but no effective in other microbial population that present in the gastrointestinal. Liu et al. studied the effect of combined of alginate-calcium and the extract from citrus wilsonii tanaka on microbiological and sensory change of white shrimps (Litopenaeus vannamei) during the storage at 4°c. They suggested that the methanol extract can be used as a new eco-friendly preservative that might use to increase the time of storage of shrimps and improve the quality.

Alfonzo et al., have been studied the effect of essential oil that extracted from lemon on the quality of salted sardines [20]. They found that the concentration of all microbial group under study were decreased during the entire period of monitoring. Also, the accumulation of histamine in sardines was decreased. On the contrast, the acceptability of sensory evaluation of sardines were observed higher than those of control due to their content of several volatile organic compounds. Fisher and Phillips reported that the citrus oils were more effective on the bacteria when we're adding to the cabbage leaf than on chicken skin [21]. Also, Tassou and Nychas found that the high effective for the essential oil were when they applied *in vitro* studies compared with in food system [22]. Furthermore, they noted that to obtain the similar effect such those *in vitro*, an increase in concentration of 10-fold, 50-fold and 25-100-fold have been used in pork sausages, soap and cheese respectively, due to that there is a large of nutrients which available in food than in culture medium, since those nutrients enabling the cells to repair themselves. Smith-Palmer et al. suggested that the concentration required on salad and vegetables is less than that on high lipid foodstuffs [23]. The effect of orange and lemon extracts in cooked Swedish style meat balls was investigated by Fernàndez-Lòpez et al. [24], the rancidity-susceptible meat products have significant advantages.

After 24 hours of storage, bologna sausage sample that have been treated with the waste water (5-10%) obtained during the dietary fiber extraction from citrus exhibited to reduce the lipid oxidation and residual nitrite levels [25]. In another study, investigated the effect of three Essential Oils (EOs) have been extracted from orange peel (*Citrus aurantium L.*), lemon peel (*Citrus limonum L.*) and bergamot peel (*Citrus aurantium L.*), as antimicrobial and antioxidant in quality of sardine pilchardus. The results showed that the growth of inoculated *S. aureus* at a level of 3.5 log₁₀ cfu/g were reduced completely from second day until the end of storage at 8°c as a result of added the EOs from bergamot citrus peel at concentration 4×0.25-0.4 μ L/ml. these result suggested that the EOs could be used as a potent natural preservative to improve the quality of sardines [26].

Milk and milk products

Kavas and Kavas, revealed that adding the orange Essential Oil (EO) to egg white protein powder as based film reduced the growth of microorganism in kashar cheese [27]. Thereby, extend the storage period than the film without EO. The phenolic compounds that present in EO inhibit the growth of pathogenic bacteria such as *S. aureus, L. monocytogenes* and *E. coli* O157:H7. In another study reported that using of *citrus aurantium L.* flowers (Bahanarang Extract (BE)) as a source of antioxidant and antimicrobial at different concentration 500 to 2000 ppm to improve the quality and extend the storage period for traditional yoghurt stew. They found that the BE had strong effect on the growth of pathogenic bacteria (i.e. *Pseudomonas aeruginosa, Escherichia coli* O157:H7, *Bacillus cereus* and *Staphylococcus aureus*). Also, decreased the protein and lipid oxidation during storage [28].

Min et al. divulge that potentially use the acid hydrolysis extract from *Citrus unshiu* peel as a natural antimicrobial for low fat, whole

Al-Dalali S

Austin Publishing Group

Table 1:									
Types of Citrus	Type of Extracts	Parts	Applied in foods	Results	References				
Citrus (kaffir lime, lime and pomelo)	Ethanol and Ethyl acetate	Peel	Stored raw chicken drumettes	More effective against pathogenic bacteria at level ranged from 0.4-50 mg/ml. The sensory properties have been increased than the control during the period of storage (14 days)	[13]				
Citrus limon	extracts	essential oil	Minced beef meat	CLEO was more effective against gram positive and negative bacteria at concentration varied from 0.039 to 1.25 mg/ml and 0.25 to 2.5 mg/ml respectively. Adding 0.06 and 0.312 mg/g from CLEO were prohibit the contamination in pathogenic bacteria, particularly <i>L.</i> <i>monocytogenes.</i> The values pf pH and TBARS were lower than those which found in control meat.	[14]				
Lemon grass and lime peel	extracts	fresh and oil	chicken patties	The optimum concentration that have been effective against TBARS and lipid oxidation was 0.25% of tested oil extracts. Reduce the pH, aerobic plate count, mould and yeast.	[15]				
Kinnow	extract	rind powder	salted chicken patties	Adding both of the kinnow rind powder extract and salt at the same time led to reduce the TBARS formation to 90%. Reducing the lipid oxidation during the storage to 39%.	[16]				
Citrus sinensis	Powder	Peel	chickens	More significantly affected on Escherichia coli in the	[18]				
Citrus wilsonii tanaka	methanol extract		white shrimps (<i>Litopenaeus</i> <i>vannamei</i>)	Increase the time of storage of shrimps. Improve the quality.	[19]				
Lemon	extract	essential oil	salted sardines	Decrease the concentration microbial groups. Decrease the accumulation of histamine. The acceptability of sensory evaluation of sardines were observed higher than those of control.	[20]				
Citrus		essential oil	pork sausages, soap and cheese	Increase the concentration of 10-fold, 50-fold and 25-100-fold that have been used in pork sausages, soap and cheese respectively, gave similar effect such those <i>in vitro</i> .	[22]				
Orange and lemon	extract		cooked Swedish style	Enhance the rancidity-susceptible meat products.	[24]				
Citrus	waste water	waste water (5-10%) obtained during the dietary fiber extraction	bologna sausage	Reduce the lipid oxidation and residual nitrite levels.	[25]				
Orange peel (<i>Citrus</i> aurantium L.), lemon peel (<i>Citrus limonum</i> L.) and bergamot peel (<i>Citrus</i> aurantium L.)	extract	essential oils	Sardine pilchardus	Adding EOs from bergamot citrus peel at concentration 4 × 0.25-0.4 μL/ml completely reduced the growth of inoculated <i>S. aureus</i> at a level of 3.5 log ₁₀ cfu/g.	[26]				
Orange	extract	essential oil	kashar cheese	Extend the storage period. Inhibit the growth of pathogenic bacteria such as <i>S. aureus, L.</i> <i>monocytogenes</i> and <i>E. coli</i> o157:H7.	[27]				
Citrus aurantium L.	Bahanarang extract (BE)	flowers	Traditional yoghurt stew	BE had strong effect on the growth of pathogenic bacteria (i.e. <i>Pseudomonas aeruginosa, Escherichia</i> <i>coli</i> o157:H7, <i>Bacillus cereus</i> and <i>Staphylococcus</i> <i>aureus</i>). Decrease the protein and lipid oxidation during storage.	[28]				
Citrus unshiu	acid hydrolysis	peel	Low fat, whole and	Potentially using the acid hydrolysis extract from citrus	[29]				
Citrus	extracts		milk	A combination of 75Mpa homogenization with either 150 ppm citrus extract or 0.12% vanillic acid, could be extend the shelf life by 4 days. Inhibit the growth all of enterobacteria. <i>Pseudomonas</i> and Lactic acid bacteria.	[30]				
Bergamot (<i>Citrus</i> bergamia), lemon (Citrus lemon) and sweet orange (<i>Citrus sinensis</i>)	extract	essential oils	cabbage leaf and chicken skin	Showed antimicrobial effective against pathogenic bacteria, i.e. Staphylococcus aureus, Escherichia coli o157:H7, Bacillus cereus, Campylobacter jejuni and Listeria monocytogenes.	[21]				
Citrus of mandarin		wastes (both of peel and leaf)	Food grade kraft paper	The Water Vapor Permeability (WVP), Peroxide Value (PV) and air transmission for the treated papers were reduced than those in original papers. Possibility using the treated paper as food packaging and food coating to protect and extend the shelf life of food from deterioration that caused by moisture and oxvoen.	[32]				
Citrus C. limon C. limonia C. latifolia and C. aurantifolia	extract	essential oils	fruit fungal diseases	The MIC was effective against <i>B. cinerea</i> , in which the extracts from <i>C. limon and C. limonia</i> displayed 312µg/ml. whereas, the extracts from <i>C. latifolia and C.</i> <i>aurantifolia</i> were 625µg/ml.	[32]				

Al-Dalali S

Austin Publishing Group

		1	1		
Lemon grass	extract	essential oils	food spoiling yeast	Using thermal and lemon grass oil together have been	[33]
				The MIC representative quality.	
Citrus (lemon, biocitro and neroli)	extracts		apple juice	The MIC ranged from 160 to 500 ppm against	[34]
				two strains forming the spores (Alicyclobacillus	
				acidoterrestris), when the biocitro and lemon extracts	
				were used.	
Citrus	lemon fruit	essential oil	apple juice	Adding 200 µL of lemon EO combined with	[35]
				temperature of 54-60°c have been shown inactivation	
				of <i>E. coli</i> O157:H7.	
Citrus	extract		pineapple juice	Using of citrus extract and sodium-benzoate reduced	[36]
				the presence of spores (Fusarium oxysporum).	
Lime		juice	honey and lime juice mixes	The antibacterial in the mixture have been affected	[37]
				against the bacteria that causes of cough (S. aureus	
				and S. pvogenes).	
Lemon juice citrus juice			pistachio nuts	The heating treatment with 15 ml of lemon, 30 ml of	[38]
		juice		water and 2.25 g of citric acid at 120°C for 1 h have	
				been decreased the AFB1 level into 49.2%.	
Citrus	extract	essential oil vapor	cucumber and lettuce	The Enterococcus faecalis and Enterococcus faecium,	[7]
				were reduced.	
Lime	extract	essential oil	pastries and cream-	MIC of the EO against the pathogenic were found 8	[39]
			filled cakes	and 4 µL/disc.	
Citrus	extract	essential oils	dates	The citrus oils coated by chitosan inhibit the incidence	[40]
				of fungal decay at 12 days of storage on the average	
				of 52-62%.	
Orange peel sweet	extract	essential oils	food contact surfaces	The essential oil has been showed inactivate for Vibrio	[44]
			(plastic cutting board	parahaemolyticus, E. coli and Salmonella typhimurium.	
			pieces and stainless	Whereas no effect on Staphylococcus aureus, on the	
			steel)	food contact surfaces	
			5.501/		

and skim milk [29]. A study was conducted by [30] to explores investigate the combination of homogenization, citrus extracts and vanillic acid, to inhibit the growth all of *enterobacteria*, *Pseudomonas* and Lactic acid bacteria. The result highlighted that a combination of 75Mpa homogenization with either 150ppm citrus extract or 0.12% vanillic acid, could be extend the shelf life by 4 days in milk.

Fruits and vegetables

The Essential Oils (EOs) that extracted from bergamot (Citrus bergamia), lemon (Citrus lemon) and sweet orange (Citrus sinensis) have been showed antimicrobial effective against pathogenic bacteria, i.e. Staphylococcus aureus, Escherichia coli o157:H7, Bacillus cereus, Campylobacter jejuni and Listeria monocytogenes. In the food system, all the assessed bacteria were less susceptible in food system (cabbage leaf and chicken skin) comparing to their susceptibility in vitro [21]. In another study was conducted by Kasaai and Moosavi [31] to investigate the citrus wastes (both of peel and leaf of mandarin), and used it to treat food grade kraft paper. The result showed that the Water Vapor Permeability (WVP), Peroxide Value (PV) and air transmission for the treated papers were reduced than those in original papers. They concluded that a possibility using the citrus waste to treat the paper used it for food packaging and food coating to protect and extend the shelf life of food from deterioration that caused by moisture and oxygen. Simas et al. [32], studied the effect of essential oils that extracted from citrus against fruit fungal diseases. The results showed that the minimum inhibitory concentration was effective against B. cinerea, in which the extracts from C. limon and C. limonia displayed 312µg/ml. whereas, the extracts from C. latifolia and C. aurantifolia were 625µg/ml. The effect of lemon grass oil against several yeast that cause food spoiling was investigated by [33]. Both of lemon grass oil alone and a mixture of all oil and thermal treatment were applied in mixed fruit juices. The result showed that using the thermal and lemon grass oil together have been established a high preservative quality than using the individual exposure.

Fruit juices

Different type of citrus extracts (lemon, biocitro and neroli) investigated to their ability for the inhibition of the two strains forming the spores (Alicyclobacillus acidoterrestris). The results showed that the MIC ranged from 160 to 500ppm when the biocitro and lemon extracts were used; additionally, the using of thermal treatment at 80°C for 3-6 min enhanced their effect in apple juice and in vitro condition [34]. In another study conducted by Espina et al. [35], to investigated effect the combined treatment of thermal treatment and citrus fruit essential oil in apple juice to inactivate of Escherichia coli O157:H7. They observed that the adding 200µL of lemon EO combined with temperature of 54-60°C in the inactivation of E. coli O157:H7, might suppose a reduction in the treatment time and temperature (5.7 min and 4.5°C) respectively. [36], stated that the using of citrus extract and sodium-benzoate reduced the presence of spores (Fusarium oxysporum) in pineapple juice and improved the effect of homogenization in the same beverage.

Honey

The stability of antibacterial activity in the honey and lime juice mixes was conducted by [37]. The heating to 100°C for 15 min did not effect in antibacterial activity of the mixture. The antibacterial in the mixture have been affected against the bacteria that causes of cough (*S. aureus* and *S. pyogenes*).

Nut

The combined effect between lemon juice, citrus juice and heating was investigated in pistachio nuts to decrease of Aflatoxin B1 (AFB1). The result showed that the heating treatment with 15ml of lemon, 30ml of water and 2.25g of citric acid at 120°C for 1 h have been decreased the AFB1 level into 49.2% matching to their level before treatment, without altering in the physical properties of pistachios nuts [38].

Salad

Fisher K et al., divulge the effect of blend of citrus essential oil vapor against Vancomycin Susceptible (VSE) and Vancomycin Resistant (VRE), i.e. *Enterococcus faecalis* and *Enterococcus faecium*, on cucumber and lettuce [7]. The result display that the VSE and VRE strains were reduced comparing to their load before expose to the essential oil vapor. Also, they advised that using essential oil vapor as alternative to chemical antibacterial.

Cake and pastries

The effect of lime Essential Oil (EO) was studied against *Staphylococcus epidermidis* and *Bacillus subtilis* whose isolated from pastries and cream-filled cakes by [39]. MIC of the EO against the pathogenic were found 8 and 4 μ L/disc. Also, they recommended that using of *citrus aurantifolia* EO in high concentration can improve the storage period for pastries and cream-filled cake.

Dates and others

The effect of different type of citrus Essential Oils (EO), combined with locust bean gum and chitosan on the Aspergillus flavus that caused postharvest spoilage in dates was conducted by [40]. They found that the citrus oils coated by chitosan inhibit the incidence of fungal decay at 12 days of storage on the average of 52-62%. [41], conducted the effect of Freeze Dried Citrus Peel (FDCP) at different levels (5.55, 11.1 and 22.2 g/kg) during the storage periods of feed, comparing to feed treated with an Antifungal Agent (AA) at 0.05g/ kg and the untreated feed. The results showed sign of adding FDCP inhibit the deterioration and infection with aflatoxin over 21 days, whereas AA treatment and untreated were deteriorated after 16 days. In another study was conducted by [42], they stated that the effect of sweet lime peel oil and orange peel oil were lower than lemon peel oil in the inhibitory effect of food borne pathogens. Also, Ramadan et al. [43] assessed the effect methanol and ethanol peel extracts (orange, lemon and pomegranate) against food borne pathogens, which has been taken from rinsate of chicken skin. The results showed that the treated of skin chicken with 5mg/ml of the extracts gave significant reduction in food borne pathogens (Figure 1) [44], investigated the effect of essential oils extracted from orange peel sweet against some type of pathogenic bacteria, which were inoculated on the food contact surfaces (plastic cutting board pieces and stainless steel). They observed that the essential oil extracted from orange showed inactivate for Vibrio parahaemolyticus, E. coli and Salmonella typhimurium. Whereas no effect on Staphylococcus aureus, on the food contact surfaces.

Conclusion

A tale sort of bundling that consolidates dinners bundling substances with antimicrobial assets to control microbial floor pollution of fixings to upgrade item microbial wellbeing and to extend timeframe of realistic usability is pulling in leisure activity in the bundling business. In concurrence with the advanced pattern of offering cost to normal and sustainable assets, the utilization of natural antimicrobial mixes, exceptionally in sustenance and biomedical applications, will turn out to be extremely visit. The immediate expansion of common mixes to sustenance is the most regular methodology of utilization, regardless of whether severa endeavors have been settled on to find decision answers for the goal of keeping away from bothersome inactivation. Plunging, showering, and covering treatment of suppers with enthusiastic alternatives are right now used to item preceding bundling as genuine choices.

References

- 1. Shahidi F, Zhong Y. Novel antioxidants in food quality preservation and health promotion. Eur J Lipid Sci Technol. 2010; 112: 930-940.
- Aziz M, Karboune S. Natural antimicrobial/antioxidant agents in meat and poultry products as well as fruits and vegetables: A review. Crit Rev Food Sci Nutr. 2017; 8398: 1-26.
- Ahmad SR, Gokulakrishnan P, Giriprasad R, Yatoo MA. Fruit-based natural antioxidants in meat and meat products: A review. Crit Rev Food Sci Nutr. 2015; 55: 1503-1513.
- Lucera A, Costa C, Conte A, Del Nobile MA. Food applications of natural antimicrobial compounds. Front Microbiol. 2012; 1-13.
- Cushnie TP, Lamb AJ. Antimicrobial activity of flavonoids. International Journal of Antimicrobial Agents. 2005; 5: 343-356.
- Chaisawadi S, Thongbute D, Methawiriyaslip W, Pitakworarat N, Chaisawadi A, Jaturonrasamee K, et al. Preliminary study of antimicrobial activities on medicinal herbs of Thai food ingredients. Acta Horticulturae 675: 2003; 675: 111-114.
- Fisher K, Phillips C, Mcwatt L. The use of an antimicrobial citrus vapour to reduce *Enterococcus sp.* on salad products. International Journal of Food Science and Technology. 2009; 44: 1748-1754.
- Mandrell RE, Wachtel MR. Novel detection techniques for human pathogens that contaminate poultry. Current Opinion in Biotechnology. 1999; 10: 273-278.
- Irkin R, Esmer OK. Novel food packaging systems with natural antimicrobial agents. J Food Sci Technol. 2015; 52: 6095-6111.
- Sánchez-ortega I, García-almendárez BE, Santos-lópez EM, Amaro-reyes A, Barboza-corona JE, Regalado C. Antimicrobial Edible Films and Coatings for Meat and Meat Products Preservation. 2014.
- Codex Alimentarius, Code of hygienic practice for meat. Codex Alimentarius Commision/Recommended Code of Practice. 58- 2005. New Zealand, FAO/ WHO, 2005.
- ICMSF. Microorganisms in Foods 6. Microbial Ecology of Food Commodities, 2nd edition. 2005.
- Klangpetch W, Phromsurin K, Hannarong K, Wichaphon J, Rungchang S. Antibacterial and antioxidant effects of tropical citrus peel extracts to improve the shelf life of raw chicken drumettes. International Food Research Journal. 2016; 23: 700-707.
- 14. Ben Hsouna A, Ben Halima N, Smaoui S, Hamdi N. Citrus lemon essential oil: chemical composition, antioxidant and antimicrobial activities with its preservative effect against Listeria monocytogenes inoculated in minced beef meat. Lipids in Health and Disease. 2017; 16: 146.
- Ibrahim HM, Salem FMA. Effect of adding lemongrass and lime peel extracts on chicken patties quality. Journal of Applied Sciences Research. 2013; 9: 5035-5047.
- Devatkal SK, Narsaiah K, Borah A. The effect of salt, extract of kinnow and pomegranate fruit by-products on colour and oxidative stability of raw chicken patties during refrigerated storage. J Food Sci Technol. 2011; 48: 472-477.
- Devatkal SK, Narsaiah K, Borah A. Anti-oxidant effect of extracts of kinnow rind, pomegranate rind and seed powders in cooked goat meat patties. Meat Sci. 2010; 85: 155-159.
- Ebrahimi A, Santini A, Alise M, Pourhossein Z, Miraalami N, Seidavi A. Effect of dried *Citrus sinensis* peel on gastrointestinal microbiota and immune system traits of broiler chickens. Italian Journal of Animal Science. 2015; 14: 712-717.
- Liu X, Jia Y, Hu Y, Xia X, Li Y, Zhou J. Effect of *Citrus wilsonii* Tanaka extract combined with alginate-calcium coating on quality maintenance of white shrimps (*Litopenaeus vannamei boone*). Food Control. 2016; 68: 83-91.

Al-Dalali S

- Alfonzo A, Martorana A, Guarrasi V, Barbera M, Gaglio R, Santulli A, et al. Effect of the lemon essential oils on the safety and sensory quality of salted sardines (*Sardina pilchardus Walbaum* 1792). Food Control. 2017; 73: 1265-1274.
- 21. Fisher K, Phillips CA. The effect of lemon, orange and bergamot essential oils and their components on the survival of *Campylobacter jejuni, Escherichia coli* 0157, *Listeria monocytogenes, Bacillus cereus* and *Staphylococcus aureus in vitro* and in food systems. J Appl Microbiol. 2006; 101: 1232-1240.
- Tassou CC, Nychas GJE. Antimicrobial activity of essential oil of mastic gum on Gram positive and Gram negative bacteria in broth and in model food system. Int Biodeterior Biodegradation. 1996; 14: 411-420.
- Smith-Palmer A, Stewart J, Fyfe L. Antimicrobial properties of plant essential oils and essences against five important food-borne pathogens. Lett Appl Microbiol. 1998; 26: 122-188.
- Fernàndez-Lòpez J, Zhi N, Aleson-Carbonell L, Pèrez-Alvarez JA, Kuri V. Antioxidant and antibacterial activities of natural extracts: applicationin beef meatballs. MeatSci. 2005; 69: 371-380.
- 25. Viuda-Martos M, Ruiz-Navajas Y, Fernández-López J, Pérez-Álvarez JA. Effect of adding citrus waste water, thyme and oregano essential oil on the chemical, physical and sensory characteristics of a bologna sausage. Innov Food Sci Emerg. Technol. 2009; 10: 655-660.
- Djenane D. Chemical Profile, Antibacterial and Antioxidant Activity of Algerian Citrus Essential Oils and Their Application in Sardina pilchardus. 2015; 208-228.
- 27. Kavas N, Kavas G. Physical-chemical and antimicrobial properties of Egg White Protein Powder films incorporated with orange essential oil on Kashar Cheese. Food Science and Technology. 2016; 36: 672-678.
- Hashemi SMB, Amininezhad R, Shirzadinezhad E, Farahani M, Yousefabad SHA. The Antimicrobial and Antioxidant Effects of *Citrus aurantium L*. Flowers (Bahar Narang) Extract in Traditional Yoghurt Stew during Refrigerated Storage. Journal of Food Safety. 2016; 36: 153-161.
- Min KY, Kim HJ, Lee KA, Kim KT, Paik HD. Antimicrobial activity of acidhydrolyzed *Citrus unshiu* peel extract in milk. Journal of Dairy Science. 2014; 97: 1955-1960.
- Bevilacqua A, D'Amato D, Sinigaglia M, Corbo MR. Combination of Homogenization, Citrus Extract and Vanillic Acid for the Inhibition of Some Spoiling and Pathogenic Bacteria Representative of Dairy Microflora. Food and Bioprocess Technology. 2013a; 6: 2048-2058.
- Kasaai MR, Moosavi A. Treatment of Kraft paper with citrus wastes for food packaging applications: Water and oxygen barrier properties improvement. Food Packaging and Shelf Life. 2017; 12: 59-65.
- 32. Simas DLR, de Amorim SHBM, Goulart FRV, Alviano CS, Alviano DS, da Silva AJR. Citrus species essential oils and their components can inhibit or stimulate fungal growth in fruit. Industrial Crops and Products. 2017; 98: 108-115.

- Tyagi AK, Gottardi D, Malik A, Guerzoni ME. Chemical composition, *in vitro* anti-yeast activity and fruit juice preservation potential of lemon grass oil. LWT-Food Science and Technology. 2014; 57: 731-737.
- Bevilacqua A, Campaniello D, Speranza B, Sinigaglia M, Corbo MR. Control of Alicyclobacillus acidoterrestris in apple juice by citrus extracts and a mild heat-treatment. 2013b; 31: 553-559.
- Espina L, Somolinos M, Ouazzou AA, Condón S, García-gonzalo D, Pagán R. Inactivation of Escherichia coli O157 : H7 in fruit juices by combined treatments of citrus fruit essential oils and heat. Int J Food Microbiol. 2012; 159: 9-16.
- Bevilacqua A, Campaniello D, Sinigaglia M, Ciccarone C, Rosaria M, Gi D. Sodium-benzoate and citrus extract increase the effect of homogenization towards spores of Fusarium oxysporum in pineapple juice. Elsevier. 2012; 28: 199-204.
- Hardoko, Yuliana F. Stability Study of Antibacterial Activity of Mixed Lime Juice and Honey of Heating Temperature on *Staphylococcus aureus* and *Streptococcus pyogenes*. International Journal of Pure and Applied Sciences and Technology. 2014; 21: 1-7.
- Rastegar H, Shoeibi S, Yazdanpanah H, Amirahmadi M, Khaneghah AM, Campagnollo FB, et al. Removal of aflatoxin B1 by roasting with lemon juice and/or citric acid in contaminated pistachio nuts. Food Control. 2017; 71: 279-284.
- 39. Jafari S, Esfahani S, Fazeli MR, Jamalifar H, Samadi M, Samadi N, et al. Antimicrobial activity of Lime essential oil against food-brone pathogens isolated from Cream-filled cakes and pastries. International journal of biological chemistry. 2011; 5: 258-265.
- 40. Aloui H, Khwaldia K, Licciardello F, Mazzaglia A, Muratore G, Hamdi M, et al. Efficacy of the combined application of chitosan and Locust Bean Gum with different citrus essential oils to control postharvest spoilage caused by Aspergillus flavus in dates. Int J Food Microbiol. 2014; 170: 21-28.
- Nam IS, Garnsworthy PC, Ahn JH. Effects of freeze-dried citrus peel on feed preservation, aflatoxin contamination and *in vitro* ruminal fermentation. Asian-Australasian Journal of Animal Sciences. 2009; 22: 674-680.
- 42. Hasija S, Ibrahim G, Wadia A. Antimicrobial Activity of *Citrus sinensis* (Orange), *Citrus limetta* (Sweet Lime) and *Citrus limon* (Lemon) Peel Oil on Selected Food Borne Pathogens. International Journal of Life Sciences Research. 2015; 3: 35-39.
- 43. Ramadan H, Min B, Tiwari AK, Reddy G, Adesiyun A, Hinton A, et al. Antibacterial activity of Pomegranate, Orange and Lemon peel extracts against food-borne pathogens and spoilage bacteria *in vitro* and on poultry skin. International Journal of Poultry Science. 2015; 14: 229-239.
- 44. Lin C, Sheu S, Hsu S, Tsai Y. Determination of bactericidal efficacy of essential oil extracted from orange peel on the food contact surfaces. 2010; 21: 1710-1715.

Ann Agric Crop Sci - Volume 4 Issue 1 - 2019 ISSN: 2573-3583 | www.austinpublishinggroup.com Al-Dalali et al. © All rights are reserved

Citation: Al-Dalali S, Zheng F and Al-Farga A. Prolonged the Shelf Life of Different Foods using the Citrus By-Products as Antimicrobials: A Review Article. Ann Agric Crop Sci. 2019; 4(1): 1039.