Clinical Image

An Unexpected Accidental Needle-Stick Injury

Watanabe T1, Kato Y2 and Baba Y3*

¹Nursing Unit of Ohu University Dental Hospital, 31-1 Misumido Tomita-machi Koriyama City, Fukushima, Japan

²Department of Oral Function and Molecular Biology, Ohu University, 31-1 Misumido Tomita-machi Koriyama City, Fukushima, Japan

³Department of General Clinical Medicine, Ohu University, 31-1 Misumido Tomita-machi Koriyama City, Fukushima, Japan

*Corresponding author: Yuh Baba, Professor, Department of General Clinical Medicine, Ohu University, Misumido Tomita-machi Koriyama City, Fukushima 963-8611, Japan

Received: March 23, 2018; Accepted: April 02, 2018;

Published: April 09, 2018

Clinical Image

A 16-year old handicapped boy was undergoing dental treatment for caries under general anesthesia in our hospital. An anesthesiologist gathered blood from him and transferred his blood into spitz tube. After the blood collection, the resident inadvertently discarded syringe mounted with an 18 gauge needle into a biological-waste plastic bag without separating the sharp waste. When the scrub nurse managed the plastic bag for biological-waste after operation, she suddenly pained in her right posterior-lateral region of the thigh. Then, she found the needle, which was mounted on blood-remaining syringe, protruded from the plastic bag, and directly realized that



Figure 1: A thick needle (18 gauge) protruded from the plastic bag.

this pain was caused by the needle-stick injury (Figure 1). Both the patient and nurse were immediately tested for HIV, Hepatitis B and Hepatitis C, resulting serologically negative. In conclusion, the health care worker must take care of not only medical practice but medical waste disposing, because needle-stick injuries sometimes occur [1].

References

 Nagao M, Iinuma Y, Igawa J, Matsumura Y, Shirano M, Matsushima A, et al. Accidental exposures to blood and body fluid in the operation room and the issue of underreporting. Am J Infect Control. 2009; 37: 541-544.