

## Research Article

# Isoproterenol has a Negative Effect on Human Submandibular Gland Cells Through B Adrenergic Receptors /Gas Signaling Pathway

Zhou T, Xu S, Huang L, Zhang J, Tian Q, Wei W\* and Wu H\*

Department of Clinical Pharmacology, Anhui Medical University, China

\*Corresponding author: Huaxun Wu and Wei Wei, Department of Clinical Pharmacology, Anhui Medical University, Key Laboratory of Anti-inflammatory and Immune Medicine, Ministry of Education, Anhui Collaborative Innovation, Tel: 13855156104, China

Received: April 12, 2022; Accepted: May 17, 2022; Published: May 24, 2022

## Abstract

Human submandibular glands play a crucial role in the function of secretion. Impairment of the glands would lead to series diseases like Sjögren's Syndrome (SS). However, mechanism of salivary glands secretion function hasn't been clarified completely. The sympathetic process may participate in the process of it as stress is one of the side effects of SS. Stress stimulates the sympathetic neuronal system causing functional changes of human submandibular glands. Therefore, our study was designed to investigate the effects of Isoproterenol (ISO) against Human Submandibular Gland Cells (HSGCs) and further explore the possible mechanisms concerned.

**Keywords:**  $\alpha$ -amylase;  $\beta$  adrenergic; Gas; Human submandibular gland cell; Isoproterenol; Sjögren's syndrome

## Abbreviations

GPCR: G Protein-Coupled Receptor; GRK2: G Protein-Coupled Receptor Kinase 2; HPA: Hypothalamic-Pituitary-Adrenal; HSG: Human Submandibular Gland; ISO: Isoproterenol; RA: Rheumatoid Arthritis; SGAG: Sulfated Glycosaminoglycans; SNS: Sympathetic Nervous System; SS: Sjögren's Syndrome  $\beta$ ; AR:  $\beta$  Adrenergic Receptors

## Introduction

Submandibular glands, parotid glands, and sublingual glands are three types of major salivary glands. They secrete lots of saliva daily, which not only contain various substances to maintain the gastrointestinal tract and oral health, but also have great significance in diseases diagnosis. Saliva plays an important role in mastication, swallowing, digestion of starch and the maintenance of teeth. The secretion of salivary glands is essential for maintaining oral health and high quality of life. Except for water and some inorganic substances like metal ions, an organic substance in saliva. There are a lot of proteins consisting of oral defense system, such as lysozyme,  $\alpha$ -amylase, mucins, lactoperoxidase, immunoglobulin, agglutinin, and others [1,2]. Dysfunction of salivary glands could lead to a lot of oral diseases, such as dental caries, dysphagia, oral mucositis and Sjögren's Syndrome (SS). SS is a prototypic chronic autoimmune disease characterized by dry mouth and dry eyes. It's mainly caused by impaired secretory function of exocrine glands such as lacrimal glands and salivary glands [3]. Damage of salivary glands can also make the electrolyte transport and the flow of saliva in the mouth stop [4-6]. Human salivary glands that are mainly composed of ductal epithelial cells, acinar cells and myoepithelial cells, play vital roles in secreting saliva.

Human salivary glands that are mainly composed of ductal epithelial cells, acinar cells and myoepithelial cells. The resting saliva is mainly produced by the submandibular gland cells [7]. Salivary

gland epithelial cells are not only the target cells of the immune response but also the effector cells of maintaining the immune response in SS [8]. Thousands of patients suffer from the salivary gland hypofunction over the world in each year. However, there are no effective pharmacological tools for the treatments of salivary dysfunction. Pathophysiology of the salivary glands damage remains to be elucidated.

It was demonstrated that prognosis of autoimmune disease patients with higher stress level is poor [9]. An elevated level of stress could lead to the activation of Hypothalamic-Pituitary-Adrenal (HPA) axis and overexcitation of the Sympathetic Nervous System (SNS) [10,11]. On the one hand, HPA axis provides a critical feedback mechanism for stress, but it is passivated in many autoimmune diseases, such as systemic lupus erythematosus and rheumatoid arthritis [12]. On the other hand, dysregulation of the autonomic nervous system could also lead to a functional somatic syndrome that is characterized by many functional symptoms [13]. The salivary gland function is associated with autonomic nervous system function. Taste and chewing induce the secretion of saliva. And secretion of saliva could also be regulated by the activation of SNS under the physiological conditions [7]. However, roles of autonomic nerve effects on human salivary glands haven't been clarified. Whether the pathophysiological mechanism of SS is associated with activation of HPA axis or overexcitation of SNS are still unclear.

$\beta$  adrenergic receptors ( $\beta$  AR) are a type of G protein-coupled receptor (GPCR) that are composed of  $\beta$ 1 AR,  $\beta$ 2 AR and  $\beta$ 3 AR and can be stimulated by autonomic nervous system or circulating catecholamine to regulate wide biological effects, such as cardiovascular activity, carbohydrate metabolism [14]. The classical signaling pathways that  $\beta$  AR participated in have played pivotal and common roles in many physiological activations. Male Wistar rats submandibular cells responded to  $\beta$  AR stimuli [15]. The activation of  $\beta$  ARs in human submandibular gland induces secretory granules

and cell membrane fusion, which in turn affects glandular secretion [16]. Increased activity of  $\beta_2$  AR/Gas signaling pathways protects myocardium from Stress-induced cardiomyopathy effects [17]. Isoproterenol (ISO), a kind of non-selective  $\beta$  AR agonist, has been widely regarded as a tool of  $\beta$  AR to verify the mechanism of  $\beta$  AR signaling in cell immune response concerned with many areas like cardiomyocytes, adipocytes and so on. Low level of the concentration would trigger off-target effects [18]. Non-Obese Diabetic (NOD) mice, a type of spontaneous SS animal model,  $\beta$  AR density appeared to decline after the stimulus of ISO [19]. C-reactive protein in the rat serum that triggered the inflammatory responses has been demonstrated the increased level after injection of ISO [20]. ISO whether could influence the cardiomyocytes and induce the cells apoptosis must have to be more discussed.

In this research, ISO was used to study the potential pathogenesis concerned with HSGCs damage. Our research aimed to explore whether  $\beta$  AR/Gas signal pathway participates in the process of HSGCs dysfunction induced by ISO and clarifies the relationship between SNS and dry or gland damage.

## Materials and Methods

### Materials

Human submandibular gland cells were purchased from Tong Pai Technology in China. Isoproterenol was obtained from Hefeng Pharmaceutical LTD (China).  $\beta_1$  AR antagonist CGP20712A (119K4600v) and  $\beta_2$  AR antagonist ICI118551 (102M4619v) were provided by Sigma company (USA). High glucose of DMEM is from Multicell. Fetal Bovine Serum (FBS) was purchased from Clerk company. Chondroitin sulfate of shark was obtained from Qingdao Green-extract Biology Science (China), and 1,9 Dimethylmethylene Blue zinc chloride double salt (DMMB) Assay (341088-1G) is from Sigma-Aldrich (USA). Annexin V-FITC/PI Apoptosis Detection Kit is from Vazyme Biotech (China). Penicillin/streptomycin (100 $\times$ ), 0.25% trypsin, goat antimouse IgG/HRP (ZB-2305), goat anti-rabbit IgG/HRP (ZB-2301), RIPA (P00138), PMSF (ST 506-2), DAPI (C1005) were all purchased from Beyotime Technology (USA). Bull Serum Albumin (A8002) was purchased from Solarbio Life Sciences in China. Anti- $\beta_1$  AR antibody (ab3442), anti- $\beta_2$  AR antibody (ab36956), anti-Gas antibody (ab83735), anti-Salivary  $\alpha$ -amylase antibody (ab201450) were all purchased from Abcam system (USA). Anti-GRK2 antibody (C-15) is from Santa Cruz Biotechnology (USA). Alexa Fluor 594-conjugated goat anti-rabbit IgG(H+L) (SA00006-4) was provided by Protein Tech Group (USA).

### Cell lines culture and treatments

HSG cells, a kind of originated from irradiated human submaxillary neoplastic intercalated duct cell lines of glands [21], were maintained in high glucose DMEM containing 10% fetal bovine serum and 1% penicillin/streptomycin. Medium was replaced 2 to 3 times a week, and 0.25% trypsin was used to change the density of the cells. ISO was used as stimuli and at the same time, two antagonists CGP20712A and ICI118551 were used for treating the cells. The concentrations of ISO, CGP20712A, and ICI118551 were all 10 $^{-6}$  mol/L. When the stimuli were exerted; all the cells were cultured with 5% High glucose of DMEM. All conditions of the cells were incubated in 37 $^{\circ}$ C 5% CO $_2$ . We stimulated the HSGCs by ISO (10 $^{-6}$ mol/L), subsequently, observe cells morphology. Two indicators, Sulfated

Glycosaminoglycans (SGAG) and  $\alpha$ amylase concerning the cells secretion function were detected. Cells apoptosis proportions were also measured.  $\beta$  Adrenergic Receptor ( $\beta$  AR) and part of associated signaling molecules were measured via western blot. The present study showed that ISO could change the morphology of HSGCs. ISO significantly increased the level of SGAG proportions in HSG cells supernatant and apparently decreased the expression of  $\alpha$ -amylase. ISO contributed to HSGCs apoptosis. Expressions of  $\beta_1$  AR,  $\beta_2$  AR and Gas all significantly decreased. GRK2 showed an up-regulated level under the exposure of ISO.

### Cells morphology observed by microscope

Cells were inoculated for 5 $\times$ 10 $^4$  per well in 6-well plate. ISO was added to the well for 48 hours treatment. Morphological changes of HSGCs were observed through microscope.

### SGAG measured by DMMB Dye-Binding Assay

Preparation of standard substance: 20 mg chondroitin sulfate of shark was weighed and dissolved in 10ml Phosphate Buffer Saline (PBS) at room temperature. Then, diluted the solution into 200, 160, 100, 80, 40, 20, 10, 5, 2.5 $\mu$ g/ml in sequence. The cells were treated for 48 hours in a 6-well plate and the supernatant was collected after centrifugation at 2000 rpm for 5 minutes. The cells were lysed on ice for 30 minutes by Radio Immunoprecipitation Assay (RIPA) with 1% Phenyl Methane Sulfonyl Fluoride (PMSF) in it to extract the protein. The solution was centrifuged at 14000 rpm at 4 $^{\circ}$ C for 10 minutes. Both the supernatant and the protein were stored at -80 $^{\circ}$ C. 10 $\mu$ l standard substance or the sample were added in 96-well plate, and then 200 $\mu$ l DMMB was aspirated into every well. The absorbance was measured by Multiskan Spectrum at 525nm wave length after 1 minute. The concentration was worked out, and the result was presented as the proportion of the supernatant in total SGAG.

### $\alpha$ -amylase level measured by laser scanning confocal microscopy

Adjusted the cells density to 2  $\times$ 10 $^5$ /ml and add 500 $\mu$ l to each well with a sterile coverslip in the 24-well plate. After 48 hours of treatment, solved the cells as followed: fixed the cells with 4% paraformaldehyde for 30 minutes; permeated the cells with 0.5% Triton X-100 for 10 minutes; block the nonspecific protein with Bull Serum Albumin(BSA) or 30 minutes; incubated the anti- $\alpha$ -amylase antibody at the ratio of 1:900 at 4 $^{\circ}$ C for a whole night; incubated the Alexa Fluor 594-conjugated goat antirabbit IgG(H+L) at the ratio of 1:150 at 37 $^{\circ}$ C for an hour and then DAPI for 8 minutes; cover the coverslip with Fluorescence quenching agent to the glass slide. Observe the protein expression under Laser Scanning Confocal Microscopy (LSCM, Leica SP8, Germany).

### Cells apoptosis proportion detected by annexin V-FITC/PI apoptosis detection kit

Briefly, inoculated the cells for 2.5 $\times$ 10 $^5$  per well in 6-well plate and the method of treatment was as mentioned. After 48 transferred the cells to ep tubes using 0.25% trypsin and added binding buffer 100 $\mu$ l, PI 5 $\mu$ l, ANNEXIN 5 $\mu$ l per tube for 10 minutes and finally added the binding buffer 200 $\mu$ l to each tube. The results were measured by CytoFLEX (Beckman Coulter, USA).

## Expressions of $\beta 1$ AR, $\beta 2$ AR, Gas, GRK2 measured by western blot

HSG cells were cultured and treated with the same condition in 6-well plate. After 48 hours of treatment, the cells were lysed, and the protein were extracted with the same methods as the depiction above. The protein was boiled with loading buffer (5 $\times$ ) at the ratio of 4:1 for 10 minutes and stored at -80°C. The methods are as followed: The protein was separated by 10% Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis (SDS-PAGE) and transferred to the Polyvinylidene Fluoride (PVDF) membranes; the membranes were blocked by 5% skim milk powder in Tween PBS; the primary antibodies (1:500) were incubated at 4°C for a whole night; the secondary antibodies were incubated at 37°C for 2 hours; the images were detected by Enhanced Chemi Luminescence (ECL) reagents after exposure of filters to films. The density of the bands was analyzed by the Microsoft (Image J, Broken Symmetry Software).

### Statistical analysis

Data were analyzed by SPSS 16.0 software and performed with GraphPad Prism 5.0 software. The results were presented as mean  $\pm$  SEM for at least 3 times experiments. All p valued less than 0.5 were considered significantly different from the control groups.

## Results

### Morphological features changed under exposure of ISO

HSG cells of Control group showed as spindly or fusiform shape and the edges of cells are clear. After treatment of ISO for 48 hours, the cells presented as polygonal shape or became shorter and the edges of the cells became indistinct (Figure 1).

### Proteoglycans metabolism of HSG cells was disordered induced by ISO

Compared with the Control group, the proportion of SGAG in ISO group increased after 48 hours treatment. Besides, the level of ICI118551 group was decreased compared to the ISO group after 48h treatment of  $\beta 2$  AR antagonist. However, there was no significance between CGP20712A and the ISO group. Proteoglycans metabolism was disordered induced by ISO. After treatment of ICI118551, the level of SGAG outside the cells decrease.

### $\alpha$ -amylase expressed in the cells decreased under the exposure of ISO

Results have confirmed the reality that ISO could lead to the

damage of HSG cells through decreasing the level of  $\alpha$ -amylase expression in the cells compared to the Control group. Compared with the ISO group,  $\beta 2$  AR antagonist ICI reversed the effect by increasing the  $\alpha$ -amylase level. But the effect of  $\beta 1$  AR antagonist CGP20712A on the cells didn't show any significance after 48 hours treatment (Figure 2).

### Levels of apoptotic HSG cells decreased after sustained induction of ISO


The results showed that the proportion of apoptotic cells induced by ISO significantly increased compared to the Control group.  $\beta 2$  AR antagonist ICI118551 significantly decreased the level of HSG cells apoptosis. It may suggest that  $\beta 2$  AR might be involved in the process of HSG cells apoptosis (Figure 3).

### Expression levels of $\beta$ AR and associated signaling molecules decreased under the suppression of ISO



After 48 hours of ISO ( $1 \times 10^{-6}$  mol/L) treatment, expression levels of both  $\beta 1$  AR and  $\beta 2$  AR declined compared to the Control group. The  $\beta 1$  AR antagonists CGP20712A and  $\beta 2$  AR antagonists ICI118551 showed opposite effects on HSG cells by increasing  $\beta$  AR expression. Meanwhile, GRK2 statistically increased in the ISO group compared to the Control group. CGP20712A and ICI118551 could both reverse the effects of ISO on HSG cells. Low expression of Gas was demonstrated in the ISO group compared to the Control group; CGP20712A and ICI118551 might increase its expression (Figure 4).

## Discussion

In this study, we analyzed the expression of the  $\beta$  Adrenergic Receptors /Gas Signaling Pathway protein. Mucins, secreted by salivary glands and distributed on the surface of the mucosa, are a type of proteoglycans and may be constituted by proteins and Sulfated Glycosaminoglycans (SGAG) [22,23]. SGAG is important in forming proteoglycans, and the metabolism has been demonstrated to be disordered in Osteoarthritis (OA) [24]. Elevated levels of SGAG in the cell supernatant suggest that the metabolism of proteoglycan in OA chondrocytes is in disorder. It also exists in saliva so that we hypothesized whether it is disordered in the damaged HSG cells. Proteoglycans metabolize into SGAG and release to the extracellular, leading to a significant increase level of SGAG extracellular. In our research, SGAG proportions of the supernatant were demonstrated an increasing level of the induction of ISO, and the effect was resisted by  $\beta 2$  AR antagonists ICI118551. It's indicated that  $\beta 2$  AR may




**Figure 1:** The morphology of the cell group supplemented with ISO was significantly changed compared with the control group, and the control group showed a long spindle shape with clear cell edges, while the ISO group had irregular cell shape and unclear edges in the ISO group.



participate in secretion function of mucins.  $\alpha$ -amylase is one of the important indicators to measure the function of salivary glands. In our research, the expression levels of  $\alpha$ -amylase were demonstrated to decrease under the effects of ISO, and the effect can also be inhibited by  $\beta$ 2AR antagonist ICI118551. The activity of  $\alpha$ -amylase was tested to be declined in SS patients and animal models which make asitia developed [25]. It may suggest that both the activity and expression level are decrease in SS. As for changes of HSG cells morphology, ISO changed the shape of the cell and made the cell edges blurred.

The intuitive result showed that HSG cell was impaired under the exposure of ISO. On the basis changes above, HSG cells apoptosis was measured. In our research, apoptotic cell proportions slightly elevated with the exposure of ISO and  $\beta$ 2 AR antagonist ICI118551 reversed the effects. As was known to us, apoptosis happens in many tissues in the balanced physiological context and diseases would develop if the cells lose the ability of apoptosis [26]. HSG cells apoptosis proportion was high under the stimulation of bleomycin probably through Akt/mTOR signaling pathway [27]. Many studies reported that deeper



**Table 1:** The proportion of SGAG (%) in the cell supernatant (mean  $\pm$  SEM, n=5 per group).

Group	SGAG (%)
Control	2.5825 $\pm$ 0.6224
ISO	5.143 $\pm$ 0.7475*
ISO+CGP20712A	4.8079 $\pm$ 1.9100
ISO+ICI118551	3.504 $\pm$ 0.2425#

mechanisms might be involved in caspases-3 protease activity and Bax/Bcl2 in cells apoptosis [28]. So, we hypothesize that  $\beta$ 2 AR may be involved in the cells apoptosis through regulating caspases-3 protease activity and Bax/Bcl-2 to play its role. The result of the increased HSG cell apoptosis levels is in accordance with the secreting functions of the impaired cells.

In our body, activation of sympathetic nerve would lead to a series of symptoms, such as rapid heart rate, vasoconstriction, bronchial smooth muscle relaxation, diuretic muscle relaxation, sphincter contraction, an elevated level of the adrenal medulla. Besides, sympathetic nerve stimulation of salivary glands would result in the production of saliva with a high concentration of protein, and parasympathetic nerves excitement would make the salivary glands produce the saliva which is poor in protein [29,30]. Those changes will happen if the organs are in the physiological condition. However, continuous stimulation of the sympathetic nerve, such as  $\beta$  AR agonist ISO, may be able to inversely act on the salivary glands by reducing the saliva. In our research, sustained effects for 48 hours of ISO in high concentration may impair the cells through decreasing the sensitivity of sympathetic nerve system. To a large extent, in the salivary glands, impairment would also happen in the same condition (Table 1).

## Conclusion

Our research provided evidence that ISO can impair the function of HSG cells.  $\beta$  AR/ $G\alpha$ s pathway may be involved in the mechanism of HSG impairment as the expression of  $\beta$ 1 AR,  $\beta$ 2 AR,  $G\alpha$ s all decreased after stimulated 48 hours by ISO. Declining functions of

human submandibular gland cells induced by isoproterenol.  $\beta$  AR /  $G\alpha$ s signal pathway participates in the impairment of HSGCs. It helps to provide a new envisage to treat many oral diseases like SS.

## Declaration of Competing Interested

W.W. and H.W. contributed to the design of the study, served as the study coordinators and edited the manuscript. T.Z, S.X designed the study, performed experiments, collected and analyzed the data and wrote the manuscript. Performed experiments and collected and analyzed the data. L.H, J Z and Q T helped perform experiments and analyzed data. All authors have read and approved the final manuscript.

## Acknowledgement

This study was supported by the National Nature Science Foundation of China (Nos 81973332 and 81302784), Provincial Natural Science Research Project of Anhui Province (KJ2018A0166), and Key Projects of Anhui Province University Outstanding Youth Talent Fund (gxyqZD2018023). Research Fund of Anhui Institute translational medicine (2021zhyx-C24).

## References

- Saruta J, To M, Sakaguchi W, Kondo Y, Tsukinoki K. Brain-derived neurotrophic factor is related to stress and chewing in saliva and salivary glands. JAPANESE DENTAL SCIENCE REVIEW. 2020; 56: 43-49.
- Mattingly A, Finley JK, Knox SM. Salivary gland development and disease. Wiley Interdiscip Rev Dev Biol. 2015; 4: 573-590.
- Chen XY, Wu HX, W Wei. Advances in the diagnosis and treatment of Sjogren's syndrome. Clin Rheumatol. 2018; 37: 1743-1749.
- Rakonczay Z Jr, Vág J, Földes A, Nagy K, Nagy Á, Hegyi P, et al. Chronic inflammation in the pancreas and salivary glands—lessons from similarities and differences in pathophysiology and treatment modalities. Curr Pharm Des. 2014; 20: 1104-1120.
- Ramaekers BL, Joore MA, Grutters JP, van den Ende P, Jong Jd, Houben R, et al. The impact of late treatment-toxicity on generic health-related quality of life in head and neck cancer patients after radiotherapy. Oral Oncol. 2011; 47: 768-774.
- Jang SI, Ong HL, Gallo A, Liu X, Illei G, Alevizos I. Establishment of functional acinar-like cultures from human salivary glands. J Dent Res. 2015; 94: 304-311.

7. Carpenter GH. The secretion, components, and properties of saliva. *Annu Rev Food Sci Technol.* 2013; 4: 267-276.
8. Campos J, Hillen MR, Barone F. Salivary Gland Pathology in Sjogren's Syndrome. *Rheum Dis Clin North Am.* 2016; 42: 473-483.
9. Capellino S, Straub RH. Neuroendocrine immune pathways in chronic arthritis. *Best Pract Res Clin Rheumatol.* 2008; 22: 285-297.
10. Quas JA, Yim IS, Oberlander TF, Nordstokke D, Essex MJ, Armstrong JM, et al. The symphonic structure of childhood stress reactivity: patterns of sympathetic, parasympathetic, and adrenocortical responses to psychological challenge. *Dev Psychopathol.* 2014; 26: 963-982.
11. Skopouli FN, Katsiogiannis S. How stress contributes to autoimmunity-lessons from Sjogren's syndrome. *FEBS Lett.* 2018; 592: 5-14.
12. Härle P, Möbius D, Carr DJ, Schölmerich J, Straub RH. An opposing time-dependent immunomodulating effect of the sympathetic nervous system conferred by altering the cytokine profile in the local lymph nodes and spleen of mice with type II collagen-induced arthritis. *Arthritis Rheum.* 2005; 52: 1305-1313.
13. Kiba T, Kanbara K, Ban I, Kato F, Kawashima S, Saka Y, et al. Saliva amylase as a measure of sympathetic change elicited by autogenic training in patients with functional somatic syndromes. *Appl Psychophysiol Biofeedback.* 2015; 40: 339-347.
14. Zheng M, Zhu W, Han Q, Xiao RP. Emerging concepts and therapeutic implications of betaadrenergic receptor subtype signaling. *Pharmacol Ther.* 2005; 108: 257-268.
15. Carpenter GH, Proctor GB, Ebersole LE, Garrett JR. Secretion of IgA by rat parotid and submandibular cells in response to autonomic stimulation *in vitro.* *Int Immunopharmacol.* 2004; 4: 1005-1014.
16. Ding C, Cong X, Zhang Y, Li SL, Wu LL, Yu GY. Beta-adrenoceptor activation increased VAMP2 and syntaxin-4 in secretory granules are involved in protein secretion of submandibular gland through the PKA/F-actin pathway. *Biosci Rep.* 2018; 38.
17. Cao X, Zhou C, Chong J, Fu L, Zhang L, Sun D, et al. Estrogen resisted stress-induced cardiomyopathy through increasing the activity of beta(2)AR-Galphas signal pathway in female rats. *Int J Cardiol.* 2015; 187: 377-386.
18. Copik AJ, Baldys A, Nguyen K, Sahdeo S, Ho H, Kosaka A, et al. Isoproterenol acts as a biased agonist of the alpha-1A-adrenoceptor that selectively activates the MAPK/ERK pathway. *PLoS One.* 2015; 10.
19. Hu Y, Purushotham KR, Wang P, Dawson R Jr, Humphreys-Beher MG. Downregulation of betaadrenergic receptors and signal transduction response in salivary glands of NOD mice. *Am J Physiol.* 1994; 266.
20. Berton G, Cordiano R, Palmieri R, Pianca S, Pagliara V, Palatini P. C-reactive protein in acute myocardial infarction: association with heart failure. *Am Heart J.* 2003; 145: 1094-1101.
21. Nelson J, Manzella K, Baker OJ. Current cell models for bioengineering a salivary gland: a minireview of emerging technologies. *Oral Dis.* 2013; 19: 236-244.
22. Çelebioğlu HY, Lee S, Chronakis IS. Chronakis, Interactions of salivary mucins and saliva with food proteins: a review. *Crit Rev Food Sci Nutr.* 2020; 60: 64-83.
23. Pinzon LM, Powers JM, O'Keefe KL, Dusevish V, Spencer P, Marshall GW. Effect of mucoprotein on the bond strength of resin composite to human dentin. *Odontology.* 2011; 99: 119-128.
24. Yu J, Cao Q, Mehra R, Laxman B, Yu J, Tomlins SA, et al. Integrative genomics analysis reveals silencing of beta-adrenergic signaling by polycomb in prostate cancer. *Cancer Cell.* 2007; 12: 419-431.
25. Khalili S, Liu Y, Sumita Y, Maria OM, Blank D, Key S, et al. Bone marrow cells are a source of undifferentiated cells to prevent Sjogren's syndrome and to preserve salivary glands function in the nonobese diabetic mice. *Int J Biochem Cell Biol.* 2010; 42: 1893-1899.
26. Xu XF, Cai BL, Guan SM, Li Y, Wu JZ, Wang Y, et al. Baicalin induces human mucoepidermoid carcinoma Mc3 cells apoptosis *in vitro* and *in vivo.* *Invest New Drugs.* 2011; 29: 637-645.
27. Cai Y, Sun R, Wang R, Ren JG, Zhang W, Zhao YF, et al. The activation of Akt/mTOR pathway by bleomycin in Epithelial-to-mesenchymal transition of human submandibular gland cells: A treatment mechanism of bleomycin for mucoceles of the salivary glands. *Biomed Pharmacother.* 2017; 90: 109-115.
28. Vutharadhi S, Jolapuram U, Kodihela LD. Nutraceutical inherent of *Spinacia oleracea* Linn. methanolic leaf extract ameliorates isoproterenol induced myocardial necrosis in male albino Wistar rats via mitigating inflammation. *Biomed Pharmacother.* 2017; 85: 239-247.
29. Garrett JR. The proper role of nerves in salivary secretion: a review. *J Dent Res.* 1987; 66: 387-397.
30. Garrett JR, Suleiman AM, Anderson LC, Proctor GB. Secretory responses in granular ducts and acini of submandibular glands *in vivo* to parasympathetic or sympathetic nerve stimulation in rats. *Cell Tissue Res.* 1991; 264: 117-126.