

## Review Article

# Pharmacopoeial Standards for Therapeutic Monoclonal Antibodies: Rituximab A Case Study

Kalaivani M\*, Goyal A, Chaudhary P and Raghuvanshi RS

Indian Pharmacopoeia Commission, Ministry of Health and Family Welfare (Govt. of India), Ghaziabad, India

**\*Corresponding author:** Kalaivani M, Senior Scientific Officer, Indian Pharmacopoeia Commission, Ministry of Health and Family Welfare, Govt. of India, Sector-23, Raj Nagar, Ghaziabad-201 002, UP, India

**Email:** kalaivani.ipc@gov.in

**Received:** April 11, 2025

**Accepted:** April 24, 2025

**Published:** April 28, 2025

## Abstract

Globally, more than 160 therapeutic monoclonal antibodies (mAbs) biosimilars are approved for marketing, including 58 in India market. The rapid growth of these therapies highlights the need for streamlined regulatory oversight and robust quality assurance to ensure their efficacy and safety. The quality of therapeutic mAbs, like all medicines, is maintained through pharmacopoeial standards and established quality control strategies. Pharmacopoeias are collections of legally required quality standards for drugs and excipients used in the manufacturing of approved drugs within a country, which must be adhered to by all who produce, distribute, or oversee these medicines. Currently, general guidance for these products is available in the United States Pharmacopoeia (USP), European Pharmacopoeia (Ph.Eur.) and Indian Pharmacopoeia (IP). However, specific pharmacopoeial monographs for mAbs are limited, for example Rituximab included in the Indian Pharmacopoeia 2022 and Infliximab concentrated solution in the Ph.Eur. in 2019 (Supplement 9.6). This article discusses the challenges and opportunities in establishing quality standards for therapeutic mAbs, using the IP 2022 monograph on Rituximab as a case example. The authors also propose a harmonized approach or collaboration among leading pharmacopoeias to develop monographs for these essential therapeutics.

**Keywords:** Rituximab; Pharmacopoeial specifications; Monoclonal antibody; Regulatory; Biotherapeutics; Quality standards

## Introduction

Therapeutic monoclonal antibodies (mAbs) have emerged as a crucial class within high-molecular-weight biopharmaceuticals, demonstrating remarkable therapeutic potential. Generally, mAbs are immunoglobulin molecules derived from a single B-cell clone, produced using recombinant DNA (rDNA) technology and hybridoma technology [1-4]. Solid tumours, haematologic malignancies, psoriatic arthritis, rheumatoid arthritis, Crohn's disease, and ankylosing spondylitis are few of the chronic and life-threatening diseases for which these mAbs have revolutionised therapy strategies [5-9]. Over the last three decades, over 160 therapeutic mAbs have gained approval as treatments from leading regulatory bodies, including the Central Drugs Standard Control Organization, CDSCO; European Medicines Agency, EMA and U.S. Food and Drug Administration, FDA [10-14].

Developing quality standards for therapeutic mAbs, which are biologically derived and structurally complex, demands specialized attention, along with sophisticated testing and controls to ensure their identity, purity, and potency. The World Health Organization (WHO) has provided foundational guidance through its Technical Report Series 822, 1992: Annex 3 [15]. Additionally, manufacturers are advised to consult other National and International guidelines which include both physicochemical and biological characterization of recombinant mAbs [16-18]. These guidelines emphasize the need to assess Critical Quality Attributes (CQA) and Key Quality Attributes (KQA) using

advanced, high-resolution analytical techniques capable of detecting subtle variations in the product. The use of these techniques is crucial for maintaining product consistency and detecting any structural or functional changes.


In the present article, the authors explore the process of developing pharmacopoeial quality standards for therapeutic mAbs illustrating the approach using Rituximab as a case study for adopting these standards in an official monograph.

## Regulatory and Pharmacopoeial Aspects of Therapeutic Mabs: Worldwide

Drug regulatory agencies and the WHO provide comprehensive guidelines for the approval of mAbs as drugs and biosimilars (Table 1) [17,19]. Additionally, the WHO offers specific guidelines for the nomenclature of therapeutic mAbs [20]. The therapeutic mAbs market is expected to increase at a compound annual growth rate (CAGR) of 11.30% from its 2020 valuation of about \$125 billion to \$494.53 billion by 2030 [20-21]. The success of therapeutic mAbs and the expiration of patents have driven the development of biosimilar therapeutics. Currently, more than 160 therapeutic mAbs have been approved for marketing worldwide [23]. Table 2 lists the therapeutic mAbs approved in the US, EU, and India. Drug regulatory authorities, including the US-FDA, EMA, CDSCO and WHO, have established comprehensive guidelines for granting marketing authorization for these products,

**Table 1:** Regulatory guidelines for marketing authorization of therapeutic mAbs in US, EU and India.

S.No	Pharmacovigilance	USA (US-FDA)	EU (GaBi online)	India (CDSCO)
1	Regulatory authority	United States Food and Drugs Administration	European Medicines Agency	Central Drugs Standards Control Organization
2	Regulatory pathways	No different pathways and/or data requirements for imported drugs vs. locally manufactured drugs		Different pathways and/or data requirements for imported drugs vs. locally manufactured drugs
4	Biosimilar Guideline- Year of Publication	2010	2005	2012
5	Nomenclature	Biological qualifier scheme	No biological qualifier scheme	
7	Applicable Guidelines	Scientific considerations in demonstrating bio-similarity to a reference product	<ul style="list-style-type: none"><li>Guideline on Similar biological medicinal products</li><li>Guideline on Similar biological medicinal products containing biotechnology-derived proteins as active substance: non-clinical and clinical issues</li><li>Similar biological medicinal products containing biotechnology-derived proteins as active substance: quality issues along with product-specific guidelines</li></ul>	<ul style="list-style-type: none"><li>CDSCO guidance for industry, 2008</li><li>Submission of Clinical Trial Application for Evaluating Safety and Efficacy</li><li>Requirement for permission of New Drug Approval</li><li>Post approval changes in Biological products: Quality, Safety and Efficacy Documents</li><li>Preparation of Quality Information for Drug Submission for New Drug Approval: Biotechnological/Biological Products</li><li>Guidelines on Similar Biologics: Regulatory Requirements for Marketing authorization in India 2016.</li></ul>
8	Number of mAbs Biosimilars approved	26	37	No approval as Biosimilar but 58 rDNA based drugs are approved as 'New Drugs'



**Figure 1:** A brief overview of general steps in monoclonal antibody development.

either as innovator drugs or biosimilars (Table 1). The rapid market expansion of these molecules highlights the critical need for robust quality assurance to ensure their efficacy and safety. Pharmacopoeias play a crucial role in improving patient access to high-quality drugs, minimizing adverse effects caused by substandard medicines, and promoting consistency in drug pricing through quality assurance [24-26]. These standards, available as public compliance documents, enable independent quality verification of a product throughout its shelf life [27]. International pharmacopoeia serves as a mandatory public standard and provides an authoritative framework for assessing the identity, strength, and purity of therapeutics [28]. Additionally, it facilitates the incorporation of harmonized testing methods as quality standards, ensuring safety and quality of medicines. The Indian

Pharmacopoeia (IP) serves as the official reference for drug and pharmaceutical standards, including biopharmaceuticals approved for in India. These standards are in accordance with the provisions of the "Drugs and Cosmetics Act, 1940, and Rules" framed under it. Notably, the current edition of IP 2022 includes monographs for rituximab and rituximab injection [29-30]. The United States Pharmacopoeia (USP) has also played a key role in establishing quality standards for biologics, contributing to its growing collection of monographs and general chapters for drugs marketed in the United States. In 2012, USP initiated efforts to outline a well-defined set of quality requirements for recombinant therapeutic mAbs. This led to the introduction of the official General Chapter <129>, titled "Analytical Procedures for Recombinant Therapeutic Monoclonal Antibodies" [31].

**Table 2:** Therapeutic mAbs and its Biosimilars approved by US-FDA, EMA and CDSCO.

US (US-FDA)		Europe (EMA)				India (CDSCO)		
Biosimilar	Year of approval	Manufacturer	Biosimilar (Brand name)	Year of approval	Manufacturer	Therapeutic mAb	Year of approval	Manufacturer/ Importers
Adalimumab Biosimilar								
Adalimumab-adaz	2018	Sandoz Inc	Hyrimoz	2018	Sandoz GmbH	Adalimumab	2017	Hetero Drugs Limited
			Hefiya					
			Halimatoz					
Hulio (adalimumab-fkjp)	2020	Sandoz Inc	Imraldi	2017	Samsung Bioepis UK Limited (SBUK)		2015	Reliance Life Sciences Private Limited
Idacio (adalimumab-aacf)	2022	Fresenius Kabi USA	Amgevita	2017	Amgen Europe B.V.		2014	Cadila Healthcare Limited
Yusimry (adalimumab-aqvh)	2021	CoherusBioSciences, Inc.	Solymbic	2017				
Abrilada (adalimumab-afzb)	2019	Pfizer Inc.	Cyltezo	2017	BoehringerIngelheim International GmbH			
Hadlima (adalimumab-bwwd)	2019	Samsung Bioepis Co., Ltd.,	Hulio	2018	Viatrix Limited			
Adalimumab-adbm	2017	BoehringerIngelheim Pharmaceuticals,Inc	Idacio	2019	Fresenius Kabi Deutschland GmbH			
			Kromeya	2019				
			Amsparity	2020	Pfizer Europe MA EEIG			
Adalimumab -atto	2016	Amgen Inc.,	Yuflyma	2021	Celltrion Healthcare Hungary Kft.			
			Hukyndra	2021	StadaArzneimittel AG			
			Libmyris	2021				
Infliximab Biosimilar								
Inflectra (Infliximab-dyyb)	2016	Celltrion, Inc.	Flixabi	2016	Samsung Bioepis UK Limited (SBUK)	Infliximab	2014	Reliance Life Sciences
Avsola (infliximab-axxq)	2019	-	-	-	-		2019	M/s Biocad India Pvt. Ltd
Infliximab-abda	2017	Samsung Bioepsis Co., Ltd.,	Remsima	2013	Celltrion Healthcare Hungary Kft.		2013	M/s Johnson & Johnson Limited import
Ixifi (Infliximab-qbtX)	2017	Pfizer Inc,	Inflectra	2013	Pfizer Europe MA EEIG			
			Zessly	2018	Sandoz GmbH			
Trastuzumab Biosimilar								
Trastuzumab-dkst	2017	Mylan GmbH	Herzuma	2018	Celltrion Healthcare Hungary Kft.	Trastuzumab	2002	Roche Scientific
			Zercepac	2021	Accord Healthcare S.L.U.		2018	Intas Pharmaceuticals Limited
			Kanjinti	2018	Amgen Europe B.V., Breda		2015	Reliance Life Sciences Private Limited
			Trazimera	2018	Pfizer Europe MA EEIG		2015	Cadila Healthcare Private Limited
			Ogivri	2018	Viatrix Limited		2018	Dr Reddy Laboratories Ltd
Herzuma (trastuzumab-pkrb)	2018	CELLTRION, Inc.	-		-		2013	Biocon Ltd
Kanjinti (trastuzumab-anns)	2019	Amgen Inc.	-				2002	Taksal pharma Private Limited
Trazimera (trastuzumab-qyyp)	2019	Pfizer Ireland Pharmaceuticals	-	-	-		2018	Biocad India Pvt Ltd
Ontruzant (trastuzumab-dttb)	2019	CELLTRION, Inc.	-	-	-		2018	Vardhman Health SpecialitisPvt Ltd
Bevacizumab Biosimilar								

Bevacizumab-awwb	2017	Amgen Inc.	Mvasi	2018	Amgen Europe B.V.	Bevacizumab		Roche Limited
							2016	Hetero Drugs Limited
							2016	Intas Pharmaceuticals Limited
Vegzelma (bevacizumab-adcd)	2022	Celltrion, Inc.	Alymsys	2021	Mabxience Research SL		2016	Reliance Life Sciences Private Limited
Alymsys (bevacizumab-maly)	2022	Amneal Pharmaceuticals LLC	Aybintio	2020	Samsung Bioepis NL B.V.	-	2017	Biocon Ltd.
			Onbevzi	2021				
Zirabev (bevacizumab-bvzr)	2019	Pfizer Inc.	Abevmy	2021	Mylan IRE Healthcare Limited	-	2017	Cadila Healthcare
			Vegzelma	2022	Celltrion Healthcare Hungary Kft.	-	2018	Dr Reddy Laboratories Ltd
			Zirabev	2019	Pfizer Europe MA EEIG	-	-	-
			Oyavas	2021	STADA Arzneimittel AG	-	-	-
			Equidacent	2020	Centus Biotherapeutics Europe Limited	-	-	-
Rituximab Biosimilar								
Riabni (rituximab-arrx)	2020	Amgen, Inc.	Truxima	2017	Celltrion Healthcare Hungary Kft.	Rituximab	2015	Hetero Drugs Limited
			Blitzima	2017				
Ruxience (rituximab-pvvr)	2019	Pfizer Ireland Pharmaceuticals Cork,	Riximyo	2017	Sandoz GmbH		2015	Reliance Life Sciences
							2013	Zenotech Laboratories
Truxima (rituximab-abbs)	2018	CELLTRION, Inc.	Ruxience	2020	Pfizer Europe MA EEIG		2013	Intas Biopharmaceuticals
							2012	Taksal Limited
							2002	Roche Scientific Limited
							2017	Biocad India Pvt Ltd
							2018	Vardhman Health SpecialitisPvt Ltd
Secukinumab Biosimilar								
						Secukinumab	2015	Novartis Limited
							2015	Sandoz Limited
Canakinumab								
						Canakinumab	2011	Novartis India Pvt Limited
Natalizumab								
						Natalizumab	2018	Eisai Pharmaceuticals India Pvt Ltd
Siltuximab								
						Siltuximab	2016	Johnson & Johnson Limited
Ofatumumab								
						Ofatumumab	2016	Novartis Healthcare Private Limited
Pembrolizumab								
						Pembrolizumab	2016	MSD pharmaceuticals Private Limited
Tocilizumab								
						Tocilizumab	2009	Taksal Limited
							2018	Roche Scientific Limited
Denosumab								

								Denosumab	2018	Intas Pharmaceuticals Ltd
									2017	Dr Reddy Laboratories Ltd
Panitumumab										
								Panitumumab	2017	Dr Reddy Laboratories Ltd
Daclizumab										
								Daclizumab	2002	Roche Scientific Co
Nimotuzumab										
								Nimotuzumab	2013	Biocon Limited
Ranibizumab										
Cimerli (ranibizumab-eqrn)	2022	CoherusBioSciences, Inc.	Byooviz	2021		Samsung Bioepis NL B.V.	Ranibizumab	2007	Novartis (I) Limited	
Byooviz (ranibizumab-nuna)	2021	Samsung Bioepis Co., Ltd.	Ximluci	2022		STADA Arzneimittel AG		2019	M/s Sandoz Private Limited	
			Ranivisio	2022		Midas Pharma GmbH		2013	Intas Biopharmaceuticals	
								2014	Alcon Laboratories (India) Pvt. Ltd.	
Omalizumab										
							Omalizumab	2015	Novo Nordisk India Pvt Ltd	
								2016	M/s Sandoz Private Limited	

The European Pharmacopoeia (Ph.Eur.) has achieved important milestone in the field of biopharmaceuticals with adoption of the monograph for Infliximab concentrated solution in European Pharmacopoeia in Ph. Eur. 9<sup>th</sup> edition, year 2019 [32-33].

## Pharmacopoeial Standards as an Quality Control of Therapeutic Mabs: Rituximab a Case Study

Globally, pharmacopoeial standards for therapeutic monoclonal antibodies (mAbs) are established through two primary approaches: 1) General Chapters/Monograph provide overarching guidelines and test methods applicable to a broad range of mAbs. 2) Specific Monographs provide tailored guidance for individual mAb products, outlining specific tests and acceptance criteria based on their unique characteristics [30]. Table 3 and 4 illustrate the availability of pharmacopoeial standards for therapeutic mAbs in various pharmacopoeias, including IP, USP, Ph.Eur., and the Int Pharm. Table 3 mainly focuses on general requirements, while Table 4 presents specific monographs for individual mAb products.

### General Guideline/General Chapter for Therapeutic Mabs

The IP provides comprehensive guidance on the development and manufacturing of therapeutic monoclonal antibodies (mAbs) [29]. This guidance specifically focuses on mAbs intended for therapeutic use, excluding those used as reagents in other drug manufacturing processes, for *in vivo* diagnostics, or for prophylactic purposes. The guidance encompasses key aspects such as general principles for mAb development, product development including cloning and cell line development, process development, analytical method development, process characterization, and analytical characterization, non-clinical and clinical studies required for mAb development, and manufacturing considerations such as large-scale manufacturing

processes, process validation, lot release testing, establishment and use of reference standards, shelf-life determination, and considerations for storage and stability. Furthermore, the IP recommends adhering to the "International Nonproprietary Names (INN) for monoclonal antibodies" guidelines issued by the World Health Organization for consistent and standardized nomenclature of mAbs [20]. Figure 1 provides a brief overview of the general steps involved in the development of therapeutic mAbs.

### Specific Pharmacopoeial Monograph for Therapeutic Mabs

Globally only two pharmacopoeias provide the quality standards of specific monoclonal antibodies despite more than 160 monoclonal antibodies approval. The IP 2022 edition has introduced specific monographs for Rituximab (Drug Substance) and Rituximab injection (Drug Product). These monographs outline quality standards for rituximab encompassing physicochemical, biological, and microbiological attributes to ensure acceptable quality of both the drug substance and the drug product. General tests include assessments of appearance, extractable volume, osmolality, pH, protein content, solubility, sub-visible particulate matter and water content as appropriate. Identification tests include biological activity, peptide mapping, capillary zone electrophoresis, and isoelectric focusing. Purity assessments encompass analysis of impurities with molecular masses differing from Rituximab (using techniques like CE-SDS and SDS-PAGE), related substances (using Size Exclusion Chromatography), charged variants (using Ion Exchange Chromatography), glycan distribution (using capillary electrophoresis with fluorescence detection), and bacterial endotoxins. Other tests include IgG-isotyping and protein content determination. Potency is determined by complement-dependent cytotoxicity assays using suitable cell lines. The potency limits are established and approved during the marketing authorization process [30].

**Table 3:** Pharmacopoeial monograph and standards for therapeutic mAbs available in various pharmacopoeias.

Standard's Name	Pharmacopoeial status					
	Indian Pharmacopoeia [IP, 2022]	US Pharmacopoeia [USP, 2024]	European Pharmacopoeia [Ph.Eur. 11.2]	British Pharmacopoeia [BP, 2022]	WHO/International Pharmacopoeia [WHO, 2022]	
General requirements/ Guidelines	Therapeutic monoclonal antibodies for human use	<129> Analytical Procedures For Recombinant Therapeutic Monoclonal Antibodies	Monoclonal Antibodies For Human Use	Monoclonal Antibodies For Human Use	3.1.1 Guidelines on evaluation of monoclonal antibodies as similar biotherapeutic products (SBPs)	
Monographs	Rituximab Rituximab injection	Not Available	Infliximab conc. solution	Infliximab conc. solution	Not Available	
Ref. standard	NA	Monoclonal IgG System Suitability	Infliximab BRP	Not available	Infliximab	
		Monoclonal IgG1, mAb001			Adalimumab	
		Monoclonal IgG1, mAb002			Bevacizumab	
		Monoclonal IgG1, mAb003	Infliximab CRS		Trastuzumab	
					Cetuximab Trastuzumab	

## Challenges in Establishing Pharmacopoeial Standards for Rituximab

Therapeutic mAbs exhibit inherent structural complexity and intrinsic heterogeneity. Notably, the Indian market boasts over 18 marketed authorizations for Rituximab, reflecting its biosimilarity to the innovator product based on comparable protein structure and function [34-35]. The development of Rituximab drug substance and

drug product monographs for the IP involved extensive input and data from domestic manufacturers and importers.

A significant challenge arose from the observed diversity in quality parameters, particularly in terms of molecular size, charge, and glycosylation patterns, among products from different manufacturers/importers. Studies have consistently highlighted these variations [36-44]. This heterogeneity presented a considerable challenge in

**Table 4:** Summary of pharmacopoeial specifications of Rituximab Monographs (IP, 2022) and Infliximab concentrated solution (Ph.Eur. 11.2.).

S.No	Pharmacopoeia requirements/ Specifications/Monograph content	Pharmacopoeial Specification		
		Rituximab (Drug Substance), (IP, 2022)	Rituximab injection (Drug Product), (IP, 2022)	Infliximab concentrated solution (Drug Substance), (Ph. Eur. 11.2)
1	Host-cell-derived proteins	NMT 100 ppm	-	Limit as approved by the competent authority
2	Host-cell- and vector-derived DNA	NMT 10 ng per dose	-	
3	Category	Anticancer	Anticancer	Monoclonal antibody (TNF alfa)
4	Description	Clear colorless to pale yellow liquid free from particles that can be observed by visual observation.	Clear to opalescent, colorless to pale yellow liquid.	Opalescent or slightly opalescent, colorless or light yellow liquid.
5	Identification	Determine by method B and any two methods from method A, C, D, E	Determine by method A, B or D and C	-
5.A	Bioassay	Complies with the biological activity as described in assay		It complies with the limits of the assay (potency)
5.B	Method	Peptide mapping by HPLC	Capillary zone electrophoresis	Peptide mapping by HPLC
	Specification	The retention time of established marker peaks should be within $\pm 0.7$ minutes of the reference solution marker peaks	Positive identity is confirmed if the difference in migration time between the main peak of the reference solution and test solution is equal or less than 0.1 minute	-the profile of the chromatogram obtained with the test solution corresponds to that of the chromatogram obtained with the reference solution; -no additional peak in the chromatogram obtained with the test solution has an area greater than 0.5 per cent of the sum of the areas of peaks 1 to 20
5.C	Sodium dodecyl sulphate–polyacrylamide gel electrophoresis	The principal band in the electropherogram obtained with test solution is similar in position and intensity to the principal band in the electropherogram obtained with reference solution.		-
5.D	Method	Capillary zone electrophoresis (CZE)	Iso Electric Focusing using capillary electrophoresis (IEF-CE)	-
	Specification	Positive identity is confirmed if the difference in migration time between the main peak of the reference solution and test solution is less than or equal to 0.1 min.	pI of principal band in reference solution is $9.3 \pm 0.2$ . In the electropherogram obtained with the <i>test solution</i> , no band other than the principal band is more intense than the principal band in the electropherogram obtained with <i>reference solution</i> .	



5.E	IEF-CE	In the electropherogram obtained with reference solution, the pI of principal is 9.3 ± 0.2. In the electropherogram obtained with test solution no band other than the principal band is more intense than the principal band in the electropherogram obtained with reference solution.		NA	-
6	Tests				
6.1	pH		6.3 to 6.7		As approved by the competent authority
6.2	Osmolality		NA	Not less than 250mosmol per kg	NA
6.3	Impurities with molecular masses differing from that of Rituximab	Method 1	Capillary Electrophoresis under reducing and non-reducing conditions		Size-exclusion chromatography
		Specification	The corrected percentage area low molecular weight impurities are not more than 10.0 per cent.	Under reducing conditions: The corrected percentage area of non-glycosylated heavy chain is Not more than 2.0 per cent Under non-reducing conditions: The corrected percentage area low molecular weight impurities is not more than 10.0 per cent	sum of all peaks other than the monomer peak: maximum 2 per cent
		Method 2	Sodium dodecyl-sulfatepolyacrylamide gel electrophoresis		-
		Specification	The band(s) observed in the test solution and reference solution match in position and intensity on visual observation.		
6.4	Related substances/ related proteins	Method	Size exclusion chromatography		Capillary electrophoresis (2.2.47) under both reducing and non-reducing conditions
		Specification	Complies with the limits approved for the particular product	The sum of the peaks with retention times less than that of the principal peak is NMT 2.0 per cent, the sum of the peaks with retention times higher than that of the principal peak is NMT 7.0 per cent and the sum of the peaks with retention times lesser and higher than that of the principal peak is not more than 9.0 per cent.	Reducing conditions: — sum of all peaks other than heavy chain and light chain: maximum 2 per cent, unless otherwise justified and authorised; Non-reducing conditions: — sum of all peaks other than the principal peak: maximum 8 per cent.
6.5	Charged variants	Method	Ion-exchange liquid chromatography		Isoelectric focusing (2.2.54): use suitable agarose gels
		Specification	Acidic variants: ≤ 30 % Main peak: Main peak ≥ 40 %,	Acidic variants ≤ 45 % Main peak ≥ 35%	— Electropherogram obtained with the test solution is similar to the electropherogram obtained with reference solution (b) and no additional bands obtained with test solution. Isoelectric points of the principal components of the test solution and reference solution (b) do not differ by more than 0.05 pI units;
6.6	Glycan distribution	Method	Capillary electrophoresis with fluorescence detection		Any suitable method as per general chapter 'Glycan analysis of glycoproteins'
		Specification	The corrected area percentage of each glycan should comply with the limits approved by National Regulatory Authority (NRA). The percent area of the peaks corresponding to galactosylated glycan should be between 35 and 65 per cent.	The corrected area percentage of each glycan should comply with the limits approved by National Regulatory Authority (NRA). The percent area of the peaks corresponding to galactosylated glycan should be between 35 and 65 per cent.	percentage of fucosylatedglycans: as authorized by the competent authority; percentage of afucosylatedglycans: as authorized by the competent authority; percentage of sialylatedglycans: as authorized by the competent authority.
6.7	Bacterial endotoxins		NMT 1.0 EU per mg or equivalent to EU per ml	NMT 1.0 EU per mg	NA
6.8	Protein A leachetes		Comply with the limits as approved by NRA	-	As approved by the competent authority

6.8	Tests stated under Parenteral Preparations		NA	Complies	NA
	Residual Protein A		NA	NA	Suitable immunochemical method based on an enzyme-linked immunosorbent assay (ELISA). <i>Limit:</i> As approved by the competent authority
6.9	Protein	Method	Ultraviolet/visible spectrophotometry		
		Specification	Not less than 90 per cent and Not more than 110 per cent of the stated amount of protein.		
7	Assay Potency	Method	Complement dependent cytotoxicity assay by using WIL2-S cells		
		Specification	Rituximab contains not less than 80 per cent and not more than 125 per cent of the stated potency		
8	Storage		Store at temperature as approved by NRA.	Store at 2 to 8° in an airtight container.	In an airtight container, under approved conditions
9	Labeling		NA	The label states (1) Content of rituximab in mg per ml (2) name of product with generic name (3) drug product (injection) in mg per ml (4) Potency; (4) storage temperature.	The label states the content in milligrams of protein per millilitre.

establishing a single, universally applicable monograph for Rituximab. However, to address this complexity, the IP monograph development process incorporated flexibility in several key areas.

Furthermore, the monograph provides flexibility in the use of alternative pharmacopoeial standards [45,46]. Acceptance criteria were defined as ranges based on approved specifications. The most suitable methods and specifications were finalized through a rigorous process involving collaboration with the National Institute of Biologicals, Noida. This collaborative effort, coupled with the robust IP monograph development process, ultimately led to the successful finalization of the Rituximab drug substance and injection monographs.

## Conclusion

Pharmacopoeial specifications are indispensable for ensuring the quality control and assurance of therapeutic mAbs. Compliance with these specifications is mandatory for manufacturers, national control laboratories, and drug regulatory authorities. Pharmacopoeial monographs provide robust analytical methods and their acceptance criteria for assessing the identity, purity, and potency of therapeutic mAbs. They also play a crucial role in identifying products that do not meet established quality standards (Not suitable quality, NSQ samples). The valuable knowledge gained from developing specific monographs for Rituximab and the "General requirements for therapeutic mAbs" within the IP will undoubtedly contribute to the development of robust pharmacopoeial standards for other therapeutic mAbs. Adherence to these standards will ensure the consistent quality, safety, and efficacy of therapeutic mAbs.

Despite the challenges encountered in developing and implementing these standards, it is crucial to recognize the immense therapeutic potential of mAbs, including biosimilars, in treating life-threatening diseases and improving global healthcare outcomes. By addressing these challenges and continuing to refine pharmacopoeial standards, we can ensure that patients worldwide have access to safe and effective mAb therapies.

## Acknowledgement

Authors acknowledge the technical support provided by IPC's Expert Working Group-Biological and rDNA products, National Institute of Biologicals, Noida, India and Indian Pharmacopoeia Commission (IPC), Ghaziabad, India.

## Author Contributions

All authors contributed to the conceptualization and development of the manuscript through collaborative meetings. AG and PC drafted the manuscript while MK critically reviewed the manuscript. All authors reviewed and provided substantial and comprehensive feedback on each draft of the manuscript. All authors read and approved the final manuscript.

## References

- Mitra S and Tomar PC. Hybridoma technology; advancements, clinical significance, and future aspects. *J Genet Eng Biotechnol*. 2021; 19: 159.
- Köhler G and Milstein C. Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature*. 1975; 256: 495-497.
- Cabilly S, Riggs AD, Pande H, Shively JE, Holmes WE, Rey M, et al. Generation of antibody activity from immunoglobulin polypeptide chains produced in *Escherichia coli*. *Proceedings of the National Academy of Sciences*. 1984; 81: 3273-3277.
- Holler PD, and Chik JK. Hybridoma Production. *Current Protocols in Molecular Biology*. 2009; 86: 11.4.1-11.4.29.
- Reichert JM, Rosensweig CJ, Faden LB, Dewitz MC. Monoclonal antibody successes in the clinic. *Nat Biotechnol*. 2005; 23: 1073-1078.
- Slivkowski MX, Mellman I. Antibody therapeutics in cancer. *Science*. 2013; 341: 1192-1198.
- Chan AC, Carter PJ. Therapeutic antibodies for autoimmunity and inflammation. *Nature Reviews Immunology*. 2010; 10: 301-316.
- Scott LJ. Infliximab: a review in Crohn's disease. *Drugs*. 2016; 76: 1347-1357.
- Nash P, Vanhoof J, Hall S, Arulmani U, Tarzynski-Potempa R, Unnebrink K, et al. Randomized Crossover Comparison of Injection Site Pain with 40 mg/0.4 or 0.8 mL Formulations of Methotrexate. *Clinical Therapeutics*. 2018; 40: 428-439.



10. Becker M, Schneider CK, Ebel B. 'Omics', biomarker discovery, and FDA's critical path. *New England Journal of Medicine*. 2006; 355: 2255-2257.
11. Hofmann F. The European regulatory environment for biosimilars. *Generics and Biosimilars Initiative Journal (GaBI Journal)*. 2013; 2: 189-193.
12. Snodin DJ, McCrossen SD. Guidelines and pharmacopoeial standards for pharmaceutical impurities: overview and critical assessment. *Regul Toxicol Pharmacol*. 2012; 63: 298-312.
13. Keitel S. Inside EDQM: the role of the pharmacopoeia in a globalized world. *Pharm Technol*. 2010; 34.
14. Niazi SK. Biosimilars: Harmonizing the Approval Guidelines. *Biologics*. 2022; 2: 171-195.
15. WHO good manufacturing practices for biological products. Annex 3. Replacement1 of Annex 1 of WHO Technical Report Series, No. 822.
16. Guidelines on Similar Biologics. Regulatory Requirements for Marketing Authorization in India. 2016.
17. World Health Organization. Guidelines on evaluation of biosimilars, Annex 2 of WHO Technical Report Series, No. 977. 2022: 1-52.
18. European Medicines Agency (EMA). Guideline on development, production, characterization and specification for monoclonal antibodies and related products. EMA/CHMP/BWP/532517/2008; 1-13.
19. World Health Organization. Guidelines for national authorities on quality assurance for biological products. 1992 Annex 2, WHO Technical Report Series No. 822.
20. World Health Organization. New INN monoclonal antibody (mAb) nomenclature scheme, INN Working Doc. 21.531. 2021: 1-3.
21. Ecker DM, Jones SD, Levine HL. The therapeutic monoclonal antibody market. *MAbs*. 2015; 7: 9-14.
22. Kaplon H, Chenoweth A, Crescioli S, et al. Antibodies to watch in 2022. *MAbs*. 2022; 14: 2014296.
23. Lyu X, Zhao Q, Hui J, Wang T, Lin M, Wang K, et al. The global landscape of approved antibody therapies, *Antibody Therapeutics*. 2022; 5: 233-257.
24. Belz S. Das Arzneibuch. Einwichtiger Pfeiler der Arzneimittelsicherheit [The pharmacopoeia. An important pillar of drug safety]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2006; 49: 1205-1211.
25. Soldi A. Pharmacopoeia as quality codex for the manufacturers. *Ann Ist Super Sanita*. 1975; 11: 269-280.
26. Grainger HS. The role of the pharmacopoeia in the control of pharmaceutical preparations. *Ann Ist Super Sanita*. 1975; 11: 305-313.
27. World Health Organization. WHO Expert Committee on Specifications for Pharmaceutical Preparations. Annex 1 Good pharmacopoeial Practices. WHO Technical Report Series No. 996, 2016.
28. The International Pharmacopoeia, 6th ed. [internet]. Geneva: World Health Organization; 2016.
29. Indian Pharmacopoeia. General Requirements. Therapeutic Monoclonal Antibodies for Human Use, IP. 2022; 3: 4576-4582.
30. Indian Pharmacopoeia. Rituximab Drug substance and Rituximab injection monographs, IP. 2022; 3: 4669-4682.
31. The United States Pharmacopoeia. General Chapters (129) Analytical Procedures for Recombinant Therapeutic Monoclonal Antibodies, 2022.
32. European Pharmacopoeia. Ninth ed. Monoclonal Antibodies For Human Use. European Pharmacopoeia. 2019; 10: 8.
33. European Pharmacopoeia. Tenth ed. Infliximab Concentrated Solution. European Pharmacopoeia. 2024; 11: 2.
34. Kaur T, Shukla BN, Yadav VK, Kulkarni MJ, Rao A. Comparison of glycoprofiles of rituximab versions licensed for sale in India and an analytical approach for quality assessment. *J Proteomics*. 2021; 244: 104267.
35. Duivelshof BL, Jiskoot W, Beck A, Veuthey JL, Guillaume D, D'Atri V. Glycosylation of biosimilars: Recent advances in analytical characterization and clinical implications. *Anal Chim Acta*. 2019; 1089: 1-18.
36. Sran KS, Sharma Y, Kaur T, Rao A. Post-translational modifications and glycoprofiling of palivizumab by UHPLC-RPLC/HILIC and mass spectrometry. *J Proteins Proteom*. 2022; 13: 95-108.
37. Edwards E, Livanos M, Krueger A, Dell A, Haslam SM, Mark Smales C, et al. Strategies to control therapeutic antibody glycosylation during bioprocessing: Synthesis and separation. *BiotechnolBioeng*. 2022; 119: 1343-1358.
38. Segu Z, Stone T, Berdugo C, Roberts A, Doud E, Li Y. A rapid method for relative quantification of N-glycans from a therapeutic monoclonal antibody during trastuzumab biosimilar development. *mAbs*. 2020; 12: 1750794.
39. Tiwold EK, Gyorgypal A, Chundawat SPS. Recent advances in biologic therapeutic N-glycan preparation techniques and analytical methods for facilitating biomanufacturing automation. *J Pharm Sci*. 2023; 112: 1485-1491.
40. Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P, et al. *Molecular Biology of the Cell*. 4<sup>th</sup> edition. New York: Garland Science; 2002. Analyzing Protein Structure and Function.
41. Aich U, Lakub J, Liu A. State-of-the-art technologies for rapid and high-throughput sample preparation and analysis of N-glycans from antibodies. *Electrophoresis*. 2016; 37: 1468-1488.
42. Aich U, Liu A, Lakub J, Mozdanzowski J, Byrne M, Shah N, et al. An Integrated Solution-Based Rapid Sample Preparation Procedure for the Analysis of N-Glycans From Therapeutic Monoclonal Antibodies. *J Pharm Sci*. 2016; 105: 1221-1232.
43. Yang X, Kim SM, Ruzanski R, Chen Y, Moses S, Ling WL, et al. Ultrafast and high-throughput N-glycan analysis for monoclonal antibodies. *mAbs*. 2016; 8: 706-717.
44. Prior S, Hufton SE, Fox B, Dougall T, Rigsby P, Bristow A. Participants of the study. International standards for monoclonal antibodies to support pre- and post-marketing product consistency: Evaluation of a candidate international standard for the bioactivities of rituximab. *MAbs*. 2018; 10: 129-142.
45. Zheng K, Bantog C, Bayer R. The impact of glycosylation on monoclonal antibody conformation and stability. *mAbs*. 2011; 3: 568-576.
46. Batra J and Rathore A. Glycosylation of Monoclonal Antibody Products: Current Status and Future Prospects. *Biotechnology progress*. 2016; 32.